



# Spin-dependent Fragmentation Functions

Rainer Jakob, University of Wuppertal



# Spin-dependent Fragmentation Functions

Rainer Jakob, University of Wuppertal

outline of the talk: (first version)

- systematic scheme for fragmentation functions (10 min)
- model calculations (10 min)
- model-independent bounds (10 min)
- processes involving fragmentation functions (10 min)
- presentation of a database on FF ( 5 min )



# Spin-dependent Fragmentation Functions

Rainer Jakob, University of Wuppertal

outline of the talk: (more realistic time schedule)

- systematic scheme for fragmentation functions (20 min)
- model calculations (20 min)
- model-independent bounds (10 min)
- processes involving fragmentation functions (20 min)
- presentation of a database on FF (10 min)



# Spin-dependent Fragmentation Functions

Rainer Jakob, University of Wuppertal

outline of the talk (... after a look at the final program)

- systematic scheme for fragmentation functions (20 min)
- model calculations (20 min)
- presentation of a database on FF (10 min)
- model-independent bounds (10 min)
- processes involving fragmentation functions (20 min)



# Spin-dependent Fragmentation Functions

Rainer Jakob, University of Wuppertal

outline of the talk (... after a look at the final program)

- systematic scheme for fragmentation functions (20 min)
- model calculations (20 min)
- presentation of a database on FF (10 min)
- model-independent bounds (10 min)
- processes involving fragmentation functions (20 min)

different aspects learned from works by Jaffe/Ji, Soper/Ralston, Mulders/Levelt/Tangerman/Boer/...



in current discussions there is a confusing large variety  
of different **fragmentation functions** around  
(particularly about the extraction of transversity)



in current discussions there is a confusing large variety  
of different **fragmentation functions** around  
(particularly about the extraction of transversity)

*normal* fragmentation function

$$D^{q \rightarrow h}(z) \text{ or } D_1(z) \text{ or } \hat{f}_1(z)$$



in current discussions there is a confusing large variety  
of different **fragmentation functions** around  
(particularly about the extraction of transversity)

“Collins” functions

$$H_1^{\perp(1)} \text{ or } \Delta\hat{D}_{H/i}$$

Collins

*normal* fragmentation function

$$D^{q \rightarrow h}(z) \text{ or } D_1(z) \text{ or } \hat{f}_1(z)$$





in current discussions there is a confusing large variety  
of different **fragmentation functions** around  
(particularly about the extraction of transversity)

“Collins” functions

$$H_1^{\perp(1)} \text{ or } \Delta\hat{D}_{H/i}$$

Collins

*normal* fragmentation function

$$D^{q \rightarrow h}(z) \text{ or } D_1(z) \text{ or } \hat{f}_1(z)$$

“transversity” fragmentation function

$$H_1(z) \text{ or } \hat{h}_1(z)$$

Ralston/Soper, Jaffe/Ji



in current discussions there is a confusing large variety  
of different **fragmentation functions** around  
(particularly about the extraction of transversity)

“Collins” functions

$$H_1^{\perp(1)} \text{ or } \Delta\hat{D}_{H/i}$$

Collins

*normal* fragmentation function

$$D^{q \rightarrow h}(z) \text{ or } D_1(z) \text{ or } \hat{f}_1(z)$$

“transversity” fragmentation function

$$H_1(z) \text{ or } \hat{h}_1(z)$$

Ralston/Soper, Jaffe/Ji


two-hadron interference FF

$$H_1^{\triangleleft(1)}(z) \text{ or } \delta q_I(z) \text{ and company}$$

Jaffe/Jin/Tang, Bianconi/Boffi/R.J./Radici



in current discussions there is a confusing large variety  
of different **fragmentation functions** around  
(particularly about the extraction of transversity)





in current discussions there is a confusing large variety  
of different **fragmentation functions** around  
(particularly about the extraction of transversity)

higher twist fragmentation functions

$\hat{e}(z)$  or  $E(z)$  and company

$\hat{h}_L(z), H_L(z), \hat{g}_T(z), G_T(z)$

Jaffe/Ji, Mulders/Tangerman/Boer/R.J.

$D^{q \rightarrow h}(z)$  or  $D_1(z)$  or

“transversity” fragmentation function

$H_1(z)$  or  $\hat{h}_1(z)$

Ralston/Soper, Jaffe/Ji

“Collins” functions

$H_1^{\perp(1)}$  or  $\Delta\hat{D}_{H/i}$

Collins

??

polarised

fragmentation function

$\Delta D_f^h(z)$  with

$h = \pi, K, \dots$  unpol.

Glück/Reya

FF

$H_1^{\triangleleft(1)}(z)$  or  $\delta q_I(z)$  and company

Jaffe/Jin/Tang, Bianconi/Boffi/R.J./Radici



in current discussions there is a confusing large variety  
of different **fragmentation functions** around  
(particularly about the extraction of transversity)

higher twist fragmentation functions

$\hat{e}(z)$  or  $E(z)$  and company

$\hat{h}_L(z), H_L(z), \hat{g}_T(z), G_T(z)$

Jaffe/Ji, Mulders/Tangerman/Boer/R.J.

$D^{q \rightarrow h}(z)$  or  $D_1(z)$  or

“transversity” fragmentation function

$H_1(z)$  or  $\hat{h}_1(z)$

spin-1 hadron FF

$D_{1LL}, D_{1LT}, D_{1TT}, G_{1LT},$

$G_{1TT}, H_{1LL}^\perp, H_{1LT}', H_{1LT}^\perp,$

$H_{1TT}', H_{1TT}^\perp$

Bacchetta/Mulders

“Collins” functions

$H_1^{\perp(1)}$  or  $\Delta\hat{D}_{H/i}$

Collins

??

polarised

fragmentation function

$\Delta D_f^h(z)$  with

$= \pi, K, \dots$  unpol.

Glück/Reya

FF

$H_1^{\triangleleft(1)}(z)$  or  $\delta q_I(z)$  and company

Jaffe/Jin/Tang, Bianconi/Boffi/R.J./Radici



in current discussions there is a confusing large variety of different **fragmentation functions** around (particularly about the extraction of transversity)

higher twist fragmentation functions

$\hat{e}(z)$  or  $E(z)$  and company

$\hat{h}_L(z), H_L(z), \hat{g}_T(z), G_T(z)$

Jaffe/Ji, Mulders/Tangerman/Boer

$D^{q \rightarrow h}(z)$

... and some more

“transversity” fragmentation f

$H_{\perp}(z)$  or  $\hat{h}_{\perp}(z)$

spin-1 hadron f

$D_{1LL}, D_{1LT}, D_{1TT}, G_{1LL},$

$G_{1TT}, H_{1LL}^{\perp}, H_{1LT}^{\perp}, H_{1LT}^{\perp},$

$H_{1TT}^{\perp}, H_{1TT}^{\perp}$

Bacchetta/Mulders

“Collins” functions

$H_1^{\perp(1)}$  or  $\Delta \hat{D}_{H/i}$

Collins

??

arised

ion function

$f_j(z)$  with

$\pi, K, \dots$  unpol.

Glück/Reya

ice FF

$H_1^{\triangleleft(1)}(z)$  or  $\delta q_I(z)$  and company

Jaffe/Jin/Tang, Bianconi/Boffi/R.J./Radici



**don't worry !**


**the number of independent fragmentation functions is limited**

**(actually, to just a few at leading twist)**

**and there is a simple systematics behind**



definition of correlation function for fragmentation process  $q \rightarrow h_1 \quad X$




Soper, Collins

$$\Delta_{ij}(k, P_1, X) = \sum_X \int \frac{d^4\xi}{(2\pi)^4} e^{ik \cdot \xi} \langle 0 | \mathcal{U}(0, \xi) \psi_i(\xi) | P_1, X \rangle \langle P_h, X | \bar{\psi}_j(0) | 0 \rangle$$





definition of correlation function for fragmentation process  $q \rightarrow h_1 \quad X$



Soper, Collins

$$\Delta_{ij}(k, P_1, X) = \sum_X \int \frac{d^4\xi}{(2\pi)^4} e^{ik \cdot \xi} \langle 0 | \mathcal{U}(0, \xi) \psi_i(\xi) | P_1, X \rangle \langle P_h, X | \bar{\psi}_j(0) | 0 \rangle$$


$\Delta$  is a  $4 \times 4$  matrix in Dirac space, and depends on the momentum vectors:

$$k, P_1,$$

and possibly spin vectors  $S$  (spin-1/2), or spin vector and tensor  $S, T$  (spin-1), etc.



definition of correlation function for fragmentation process  $q \rightarrow h_1 h_2 X$



Soper, Collins

$$\Delta_{ij}(k, P_1, P_2) = \sum_X \int \frac{d^4\xi}{(2\pi)^4} e^{ik \cdot \xi} \langle 0 | \mathcal{U}(0, \xi) \psi_i(\xi) | P_1, P_2; X \rangle \langle P_h, P_2; X | \bar{\psi}_j(0) | 0 \rangle$$

$\Delta$  is a  $4 \times 4$  matrix in Dirac space, and depends on the momentum vectors:

$$k, P_1, P_2$$

and possibly spin vectors  $S$  (spin-1/2), or spin vector and tensor  $S, T$  (spin-1), etc.



⇒ construct most general ansatz

Soper/Ralston, Mulders/Tangerman

for instance for one spin-0 hadron in the final state:

$$\Delta(k, P_h) = B_1 M_h + B_2 \not{P}_h + B_3 \not{k} + (B_4/M_h) \sigma_{\mu\nu} P_h^\mu k^\nu$$

with  $B_i = B_i(P_h \cdot k, k^2)$



⇒ construct most general ansatz

Soper/Ralston, Mulders/Tangerman

for instance for one **spin-0** hadron in the final state:

$$\Delta(k, P_h) = B_1 M_h + B_2 \not{P}_h + B_3 \not{k} + (B_4/M_h) \sigma_{\mu\nu} P_h^\mu k^\nu$$

with  $B_i = B_i(P_h \cdot k, k^2)$

only terms which are in accordance with constraints from:

**1.) hermiticity      2.) behavior under parity transformation**

note:  $B_4$  is T-odd, i.e. it would be forbidden by a constraint from the behavior of Dirac fields under time-reversal, if there were no final state interactions



⇒ construct most general ansatz

Soper/Ralston, Mulders/Tangerman

for instance for one spin-1/2 hadron in the final state:

$$\begin{aligned}
 \Delta(k, P_h, S) = & B_1 M_h + B_2 \not{P}_h + B_3 \not{k} + (B_4/M_h) \sigma_{\mu\nu} P_h^\mu k^\nu \\
 & + i B_5 (k \cdot S) \gamma_5 + B_6 M_h \not{S} \gamma_5 + (B_7/M_h) (k \cdot S) \not{P}_h \gamma_5 \\
 & + (B_8/M_h) (k \cdot S) \not{k} \gamma_5 + i B_9 \sigma_{\mu\nu} \gamma_5 S^\mu P_h^\nu \\
 & + i B_{10} \sigma_{\mu\nu} \gamma_5 S^\mu k^\nu + i (B_{11}/M_h^2) (k \cdot S) \sigma_{\mu\nu} \gamma_5 k^\mu P_h^\nu \\
 & + (B_{12}/M_h) \epsilon_{\mu\nu\rho\sigma} \gamma^\mu P_h^\nu k^\rho S^\sigma
 \end{aligned}$$

with  $B_i = B_i(P_h \cdot k, k^2)$

only terms which are in accordance with constraints from:

**1.) hermiticity      2.) behavior under parity transformation**

note:  $B_4, B_5$ , and  $B_{12}$  are T-odd, i.e. they would be forbidden by a constraint from the behavior of Dirac fields under time-reversal, if there were no final state interactions



fragmentation functions are obtained from the correlation function  $\Delta$   
 by projection with Dirac matrices  $\Gamma$ ,  
 and integration over components of the quark momentum

$$\Delta^{[\Gamma]}(z) \equiv \frac{1}{4z} \int dk^+ \int d^2\mathbf{k}_T \text{Tr} [\Delta\Gamma] \Big|_{k^- = P_h^- / z}$$

or

$$\Delta^{[\Gamma]}(z, -z\mathbf{k}_T) \equiv \frac{1}{4z} \int dk^+ \text{Tr} [\Delta\Gamma] \Big|_{k^- = P_h^- / z ; \mathbf{k}_T}$$

with the Dirac matrices  $\Gamma \in \{\gamma^-, \gamma^-\gamma_5, i\sigma^{\alpha-}\gamma_5, \dots, \text{higher twist}\}$



fragmentation functions are obtained from the correlation function  $\Delta$   
 by projection with Dirac matrices  $\Gamma$ ,  
 and integration over components of the quark momentum

$$\Delta^{[\Gamma]}(z) \equiv \frac{1}{4z} \int dk^+ \int d^2\mathbf{k}_T \text{Tr} [\Delta\Gamma] \Big|_{k^- = P_h^- / z ; \mathbf{k}_T}$$

or

$$\Delta^{[\Gamma]}(z, -z\mathbf{k}_T) \equiv \frac{1}{4z} \int dk^+ \text{Tr} [\Delta\Gamma] \Big|_{k^- = P_h^- / z ; \mathbf{k}_T}$$

with the Dirac matrices  $\Gamma \in \{\gamma^-, \gamma^-\gamma_5, i\sigma^{\alpha-}\gamma_5, \dots, \text{higher twist}\}$

interpretation in the context of LC quantisation:  $\Gamma$  determines quark spin states



fragmentation functions are obtained from the correlation function  $\Delta$   
by projection with Dirac matrices  $\Gamma$ ,  
and integration over components of the quark momentum

$$\Delta^{[\Gamma]}(z) \equiv \frac{1}{4z} \int dk^+ \int d^2\mathbf{k}_T \text{Tr} [\Delta\Gamma] \Big|_{k^- = P_h^- / z ; \mathbf{k}_T}$$

or

$$\Delta^{[\Gamma]}(z, -z\mathbf{k}_T) \equiv \frac{1}{4z} \int dk^+ \text{Tr} [\Delta\Gamma] \Big|_{k^- = P_h^- / z ; \mathbf{k}_T}$$

with the Dirac matrices  $\Gamma \in \{\gamma^-, \gamma^-\gamma_5, i\sigma^{\alpha-}\gamma_5, \dots, \text{higher twist}\}$

interpretation in the context of LC quantisation:  $\Gamma$  determines quark spin states

$$[\gamma^-]: \bullet \rightarrow + \leftarrow \bullet$$





fragmentation functions are obtained from the correlation function  $\Delta$   
by projection with Dirac matrices  $\Gamma$ ,  
and integration over components of the quark momentum

$$\Delta^{[\Gamma]}(z) \equiv \frac{1}{4z} \int dk^+ \int d^2\mathbf{k}_T \text{Tr} [\Delta\Gamma] \Big|_{k^- = P_h^- / z ; \mathbf{k}_T}$$

or

$$\Delta^{[\Gamma]}(z, -z\mathbf{k}_T) \equiv \frac{1}{4z} \int dk^+ \text{Tr} [\Delta\Gamma] \Big|_{k^- = P_h^- / z ; \mathbf{k}_T}$$

with the Dirac matrices  $\Gamma \in \{\gamma^-, \gamma^-\gamma_5, i\sigma^{\alpha-}\gamma_5, \dots, \text{higher twist}\}$

interpretation in the context of LC quantisation:  $\Gamma$  determines quark spin states

$$[\gamma^-]: \bullet \rightarrow + \leftarrow \bullet$$

$$[\gamma^-\gamma_5]: \bullet \rightarrow - \leftarrow \bullet$$



fragmentation functions are obtained from the correlation function  $\Delta$   
by projection with Dirac matrices  $\Gamma$ ,  
and integration over components of the quark momentum




$$\Delta^{[\Gamma]}(z) \equiv \frac{1}{4z} \int dk^+ \int d^2\mathbf{k}_T \text{Tr} [\Delta\Gamma] \Big|_{k^- = P_h^- / z ; \mathbf{k}_T}$$

or

$$\Delta^{[\Gamma]}(z, -z\mathbf{k}_T) \equiv \frac{1}{4z} \int dk^+ \text{Tr} [\Delta\Gamma] \Big|_{k^- = P_h^- / z ; \mathbf{k}_T}$$

with the Dirac matrices  $\Gamma \in \{\gamma^-, \gamma^-\gamma_5, i\sigma^{\alpha-}\gamma_5, \dots, \text{higher twist}\}$

interpretation in the context of LC quantisation:  $\Gamma$  determines quark spin states






## interpretation

Kogut, Soper, Jaffe


“good” components of Dirac field in LC quantisation

$$\psi_{\pm} = P_{\pm} \psi = \frac{1}{2} \gamma^{\mp} \gamma^{\pm} \psi$$


helicity / chiral projectors  $P_{R/L} = (1 \pm \gamma_5)/2$



$$\begin{aligned} \bar{\psi} \gamma^+ \psi &= \sqrt{2} \psi_+^\dagger (P_R P_R + P_L P_L) \psi_+ \\ &= \bar{R}R + \bar{L}L \end{aligned}$$



$$\begin{aligned} \bar{\psi} \gamma^+ \gamma_5 \psi &= \sqrt{2} \psi_+^\dagger (P_R P_R - P_L P_L) \psi_+ \\ &= \bar{R}R - \bar{L}L \end{aligned}$$



$$\begin{aligned} \bar{\psi} i\sigma^{i+} \gamma_5 \psi &= \sqrt{2} \psi_+^\dagger (P_L \gamma^i P_R - P_R \gamma^i P_L) \psi_+ \\ &= \bar{L}R - \bar{R}L \end{aligned}$$



**independent fragmentation functions at leading twist for**

**one unpolarised hadron**

**observed in a quark-jet**



$$\Delta^{[\gamma^-]}(z) = D_1 \quad \left( \bullet \rightarrow \circ \right)$$

$$\Delta^{[\gamma^- \gamma_5]}(z) = 0$$

$$\Delta^{[i\sigma^{i-} \gamma_5]}(z) = 0$$



$$\Delta^{[\gamma^-]}(z, \mathbf{k}_T) = D_1 \quad \left( \bullet \rightarrow \circ \right)$$

$$\Delta^{[\gamma^- \gamma_5]}(z, \mathbf{k}_T) = 0$$

$$\Delta^{[i\sigma^{i-} \gamma_5]}(z, \mathbf{k}_T) = 0 + \frac{\epsilon_T^{ij} k_{Tj}}{M_h} H_1^\perp$$

$$\left( \uparrow \bullet \rightarrow \circ \right) - \left( \downarrow \bullet \rightarrow \circ \right)$$




$$\Delta^{[\gamma^-]}(z, \mathbf{k}_T) = D_1 \quad (\bullet \rightarrow \circ)$$

$$\Delta^{[\gamma^- \gamma_5]}(z, \mathbf{k}_T) = 0$$

T-odd FF:  $H_1^\perp$  (Collins)  
transv. pol. quark  $\rightarrow$  unpol. hadron

$$\Delta^{[i\sigma^{i-}\gamma_5]}(z, \mathbf{k}_T) = 0 + \frac{\epsilon_T^{ij} k_{Tj}}{M_h} \mathbf{H}_1^\perp$$






remark:

$$\Delta^{[\gamma^-]}(z, \mathbf{k}_T) = D_1 \quad (\bullet \rightarrow \circ)$$

$\Delta D$  is forbidden by parity constraint

long. pol. quark  $\rightarrow$  unpol. hadron

$$\Delta^{[\gamma^- \gamma_5]}(z, \mathbf{k}_T)$$



~~$\Delta D_f^h$~~

$$\Delta^{[i\sigma^{i-}\gamma_5]}(z, \mathbf{k}_T) = 0 + \frac{\epsilon_T^{ij} k_{Tj}}{M_h} H_f^h$$

$$(\uparrow \bullet \rightarrow \circ) - (\downarrow \bullet \rightarrow \circ)$$







**independent fragmentation functions at leading twist for**


**one spin-1/2 hadron**

**observed in a quark-jet**



$$\Delta^{[\gamma^-]}(z) = D_1$$


$$\Delta^{[\gamma^- \gamma_5]}(z) = \lambda_h G_1$$


$$\Delta^{[i\sigma^{i-} \gamma_5]}(z) = S_{hT}^i H_1$$





$$\Delta^{[\gamma^-]}(z) = D_1$$

$$\Delta^{[\gamma^- \gamma_5]}(z) = \lambda_h G_1$$


$$\Delta^{[i\sigma^{i-} \gamma_5]}(z) = S_{hT}^i H_1$$


unpolarised FF  
probability a quark fragments into  
hadron  $h$  plus rest  $X$



$$\Delta^{[\gamma^-]}(z) = D_1$$


$$\Delta^{[\gamma^- \gamma_5]}(z) = \lambda_h G_1$$





$$\Delta^{[i\sigma^{i-} \gamma_5]}(z) = S_{hT}^i H_1$$


unpolarised FF  
probability a quark fragments into  
hadron  $h$  plus rest  $X$


preference in  
**longitudinal** spin transfer



$$\Delta^{[\gamma^-]}(z) = D_1$$


$$\Delta^{[\gamma^- \gamma_5]}(z) = \lambda_h G_1$$


$$\Delta^{[i\sigma^{i-} \gamma_5]}(z) = S_{hT}^i H_1$$





unpolarised FF  
probability a quark fragments into  
hadron  $h$  plus rest  $X$


preference in  
**longitudinal** spin transfer

preference in  
**transverse** spin transfer



$$\Delta^{[\gamma^-]}(z) = D_1$$


$$\Delta^{[\gamma^- \gamma_5]}(z) = \lambda_h G_1$$


$$\Delta^{[i\sigma^{i-} \gamma_5]}(z) = S_{hT}^i H_1$$


unpolarised FF  
probability a quark fragments into  
hadron  $h$  plus rest  $X$

preference in  
**longitudinal** spin transfer

preference in  
**transverse** spin transfer

the kind of information we are after to understand the process of hadronization:

... how much of the quark spin state is *remembered*  
by a leading hadron after fragmentation ?



$$\Delta^{[\gamma^-]}(z) = D_1$$

$$\left( \bullet \rightarrow \text{pink circle} \right)$$

$$\Delta^{[\gamma^- \gamma_5]}(z) = \lambda_h G_1$$

$$\left( \bullet \rightarrow \text{pink circle} \right) - \left( \bullet \rightarrow \text{pink circle} \right)$$

$$\Delta^{[i\sigma^{i-} \gamma_5]}(z) = S_{hT}^i H_1$$

$$\left( \uparrow \bullet \rightarrow \text{pink circle} \right) - \left( \downarrow \bullet \rightarrow \text{pink circle} \right)$$



Mulders/Tangerman

$$\Delta^{[\gamma^-]}(z, \mathbf{k}_T) = D_1 + \frac{\epsilon_{Tij} k_T^i S_{hT}^j}{M_h} D_{1T}^\perp$$

$$\Delta^{[\gamma^- \gamma_5]}(z, \mathbf{k}_T) = \lambda_h G_{1L} + \frac{\mathbf{k}_T \cdot \mathbf{S}_{hT}}{M_h} G_{1T}$$


$$\Delta^{[i\sigma^{i-} \gamma_5]}(z, \mathbf{k}_T) = S_{hT}^i H_{1T} + \frac{\epsilon_T^{ij} k_{Tj}}{M_h} H_{1T}^\perp$$






$$\Delta^{[\gamma^-]}(z, \mathbf{k}_T) = D_1 + \frac{\epsilon_{Tij} k_T^i S_{hT}^j}{M_h} \mathbf{D}_{1T}^\perp$$

Mulders/Tangerman




$$\Delta^{[\gamma^- \gamma_5]}(z, \mathbf{k}_T) = \lambda_1 C_1 + \frac{G_{1T}}{M_h}$$

**2 T-odd FF**  
 $D_{1T}^\perp$  and  $H_{1T}^\perp$  (Collins)  
 unpol. quark  $\rightarrow$  transv. pol. hadron  
 and  
 transv. pol. quark  $\rightarrow$  unpol. hadron



$$+ \frac{\epsilon_T^{ij} k_{Tj}}{M_h} \mathbf{H}_1^\perp$$



$$+ \frac{\lambda_h}{M_h} (\Pi_{1L} + \Pi_{1T})$$





Mulders/Tangeman

**3 "unexpected" FF**


$G_{1T}, H_{1L}^\perp$  and  $H_{1T}^\perp$

long. pol. quark  $\rightarrow$  transv. pol. hadron



and

transv. pol. quark  $\rightarrow$  long. pol. hadron



$$\frac{\epsilon_{Tij} k_T^i S_{hT}^j}{M_h} D_{1T}^\perp$$





$$+ \frac{\mathbf{k}_T \cdot \mathbf{S}_{hT}}{M_h} G_{1T}$$



$$\Delta^{[i\sigma^{i-}\gamma_5]}(z, \mathbf{k}_T) = S_{hT}^i H_{1T} + \frac{\mathbf{k}_T \cdot \mathbf{S}_{hT}}{M_h} H_{1T}^\perp$$



$$+ \frac{k_T^i}{M_h} \left( \lambda_h H_{1L}^\perp + \frac{\mathbf{k}_T \cdot \mathbf{S}_{hT}}{M_h} H_{1T}^\perp \right)$$





**independent fragmentation functions at leading twist for**

**two unpolarised hadrons**

**observed in the same quark-jet**

two hadron (interference) fragmentation functions



$$(R \equiv P_1 - P_2)$$

$$\Delta^{[\gamma^-]}(z_1, z_2, \mathbf{k}_T, \mathbf{R}_T) = D_1 \left( \bullet \rightarrow \begin{array}{c} \circ \\ \circ \end{array} \right)$$

$$\Delta^{[\gamma^- \gamma_5]}(z_1, z_2, \mathbf{k}_T, \mathbf{R}_T) = \frac{\epsilon_T^{ij} R_{Ti} k_{Tj}}{M_1 M_2} G_1^\perp$$

$$\left( \begin{array}{c} \bullet \\ \bullet \end{array} \rightarrow \begin{array}{c} \circ \\ \circ \end{array} \right) - \left( \begin{array}{c} \bullet \\ \bullet \end{array} \leftarrow \begin{array}{c} \circ \\ \circ \end{array} \right)$$

$$\Delta^{[i\sigma^{i-} \gamma_5]}(z_1, z_2, \mathbf{k}_T, \mathbf{R}_T) = \frac{\epsilon_T^{ij} R_{Tj}}{M_1 + M_2} H_1^\triangleleft + \frac{\epsilon_T^{ij} k_{Tj}}{M_1 + M_2} H_1^\perp$$

$$\left( \begin{array}{c} \bullet \\ \bullet \end{array} \rightarrow \begin{array}{c} \circ \\ \circ \end{array} \right) - \left( \begin{array}{c} \bullet \\ \bullet \end{array} \leftarrow \begin{array}{c} \circ \\ \circ \end{array} \right)$$

Bianconi/Boffi/R.J./Radici



$$(R \equiv P_1 - P_2)$$


$$\Delta^{[\gamma^-]}(z_1, z_2, \mathbf{k}_T, \mathbf{R}_T) = D_1 \quad \left( \bullet \rightarrow \begin{matrix} \circ \\ \circ \end{matrix} \right)$$

**T-odd FF**

$$G_1^\perp$$

long. pol. quark  $\rightarrow$  two hadrons

$$\Delta^{[\gamma^- \gamma_5]}(z_1, z_2, \mathbf{k}_T, \mathbf{R}_T) = \frac{\epsilon_T^{ij} R_{Ti} k_{Tj}}{M_1 M_2} G_1^\perp$$



$$\Delta^{[i\sigma^i - \gamma_5]}(z_1, z_2, \mathbf{k}_T, \mathbf{R}_T) = \frac{\epsilon_T^{ij} R_{Tj}}{M_1 + M_2} H_1^\triangleleft + \frac{\epsilon_T^{ij} R_{Ti}}{M_1 + M_2} H_1^\perp$$

$$\left( \begin{matrix} \bullet \\ \uparrow \end{matrix} \rightarrow \begin{matrix} \circ \\ \circ \end{matrix} \right) - \left( \begin{matrix} \bullet \\ \downarrow \end{matrix} \rightarrow \begin{matrix} \circ \\ \circ \end{matrix} \right)$$



$$(R \equiv P_1 - P_2)$$

$$\Delta^{[\gamma^-]}(z_1, z_2, \mathbf{k}_T, \mathbf{R}_T) = D_1 \left( \bullet \rightarrow \begin{matrix} \circ \\ \circ \end{matrix} \right)$$

**T-odd FF**


$$H_1^\triangleleft \text{ and } H_1^\perp$$

transv. pol. quark  $\rightarrow$  two hadrons two  
'variants' of the Collins function  
come with  $\mathbf{R}_T$  and  $\mathbf{k}_T$

$$= \frac{\epsilon_T^{ij} R_{Ti} k_{Tj}}{M_1 M_2} G_1^\perp$$

$$\left( \begin{matrix} \bullet \\ \blacktriangleright \end{matrix} \rightarrow \begin{matrix} \circ \\ \circ \end{matrix} \right) - \left( \begin{matrix} \bullet \\ \blacktriangleleft \end{matrix} \rightarrow \begin{matrix} \circ \\ \circ \end{matrix} \right)$$

$$\Delta^{[i\sigma^{i-}\gamma_5]}(z_1, z_2, \mathbf{k}_T, \mathbf{R}_T) = \frac{\epsilon_T^{ij} R_{Tj}}{M_1 + M_2} \mathbf{H}_1^\triangleleft + \frac{\epsilon_T^{ij} k_{Tj}}{M_1 + M_2} \mathbf{H}_1^\perp$$





**independent fragmentation functions at leading twist for**

**one spin-1 hadron**

**observed in a quark-jet**



$$\Delta^{[\gamma^-]}(z, \mathbf{k}_T) = D_1 + \frac{\epsilon_{Tij} k_T^i S_{hT}^j}{M_h} D_{1T}^\perp$$

$$\Delta^{[\gamma^- \gamma_5]}(z, \mathbf{k}_T) = \lambda_h G_{1L} + \frac{\mathbf{k}_T \cdot \mathbf{S}_{hT}}{M_h} G_{1T}$$

$$\Delta^{[i\sigma^{i-} \gamma_5]}(z, \mathbf{k}_T) = S_{hT}^i H_{1T} + \frac{\epsilon_T^{ij} k_{Tj}}{M_h} H_{1T}^\perp$$

$$+ \frac{k_T^i}{M_h} \left( \lambda_h H_{1L}^\perp + \frac{\mathbf{k}_T \cdot \mathbf{S}_{hT}}{M_h} H_{1T}^\perp \right)$$





Bacchetta/Mulders

$$\Delta^{[\gamma^-]}(z, \mathbf{k}_T) = D_1 + \frac{\epsilon_{Tij} k_T^i S_{hT}^j}{M_h} D_{1T}^\perp$$

$$+ \epsilon_T^{\mu\nu} S_{T\nu} \frac{k_{T\mu}}{M} D_{1T}^\perp + S_{LL} D_{1LL} + \frac{\mathbf{S}_{LT} \cdot \mathbf{k}_T}{M} D_{1LT} + \frac{\mathbf{k}_T \cdot \mathbf{S}_{TT} \cdot \mathbf{k}_T}{M^2} D_{1TT}$$

$$\Delta^{[\gamma^- \gamma_5]}(z, \mathbf{k}_T) = \lambda_h G_{1L} + \frac{\mathbf{k}_T \cdot \mathbf{S}_{hT}}{M_h} G_{1T}$$

$$+ \epsilon_T^{\mu\nu} S_{LT\nu} \frac{k_{T\mu}}{M} G_{1LT} + -\epsilon_T^{\mu\nu} S_{TT\nu\rho} \frac{k_T^\rho k_{T\mu}}{M^2} G_{1TT}$$

$$\Delta^{[i\sigma^{i-} \gamma_5]}(z, \mathbf{k}_T) = S_{hT}^i H_{1T} + \frac{\epsilon_T^{ij} k_{Tj}}{M_h} H_1^\perp$$

$$+ S_{LL} \frac{\epsilon_T^{ij} k_{Tj}}{M} H_{1LL}^\perp + \epsilon_T^{ij} S_{LTj} H'_{1LT} + \frac{\mathbf{S}_{LT} \cdot \mathbf{k}_T}{M} \frac{\epsilon_T^{ij} k_{Tj}}{M} H_{1LT}^\perp$$

$$+ \frac{k_T^i}{M_h} \left( \lambda_h H_{1L}^\perp + \frac{\mathbf{k}_T \cdot \mathbf{S}_{hT}}{M_h} H_{1T}^\perp \right)$$

$$+ \epsilon_T^{ij} S_{TTjl} \frac{k_T^l}{M} H'_{1TT} + \frac{\mathbf{k}_T \cdot \mathbf{S}_{TT} \cdot \mathbf{k}_T}{M^2} \frac{\epsilon_T^{ij} k_{Tj}}{M} H_{1TT}^\perp$$



Bacchetta/Mulders

$$\Delta^{[\gamma^-]}(z, \mathbf{k}_T) = D_1 + \frac{\epsilon_{Tij} k_T^i S_{hT}^j}{M_h} D_{1T}^\perp$$

$$+ \epsilon_T^{\mu\nu} S_{T\nu} \frac{k_{T\mu}}{M} D_{1T}^\perp + S_{LL} D_{1LL} + \frac{\mathbf{S}_{LT} \cdot \mathbf{k}_T}{M} D_{1LT} + \frac{\mathbf{k}_T \cdot \mathbf{S}_{TT} \cdot \mathbf{k}_T}{M^2} D_{1TT}$$

$$\Delta^{[\gamma^- \gamma_5]}(z, \mathbf{k}_T) = \lambda_h G_{1L} + \frac{\mathbf{k}_T \cdot \mathbf{S}_{hT}}{M_h} G_{1T}$$

$$+ \epsilon_T^{\mu\nu} S_{LT\nu} \frac{k_{T\mu}}{M} G_{1LT} + -\epsilon_T^{\mu\nu} S_{TT\nu\rho} \frac{k_T^\rho k_{T\mu}}{M^2} G_{1TT}$$

$$\Delta^{[i\sigma^{i-} \gamma_5]}(z, \mathbf{k}_T) = S_{hT}^i H_{1T} + \frac{\epsilon_T^{ij} k_{Tj}}{M_h} H_1^\perp$$

$$+ S_{LL} \frac{\epsilon_T^{ij} k_{Tj}}{M} H_{1LL}^\perp + \epsilon_T^{ij} S_{LTj} H'_{1LT} + \frac{\mathbf{S}_{LT} \cdot \mathbf{k}_T}{M} \frac{\epsilon_T^{ij} k_{Tj}}{M} H_{1LT}^\perp$$

$$+ \frac{k_T^i}{M_h} \left( \lambda_h H_{1L}^\perp + \frac{\mathbf{k}_T \cdot \mathbf{S}_{hT}}{M_h} H_{1T}^\perp \right)$$

$$+ \epsilon_T^{ij} S_{TTjl} \frac{k_T^l}{M} H'_{1TT} + \frac{\mathbf{k}_T \cdot \mathbf{S}_{TT} \cdot \mathbf{k}_T}{M^2} \frac{\epsilon_T^{ij} k_{Tj}}{M} H_{1TT}^\perp$$

relations between spin-1 FF

and two spin-0 FF

Bacchetta/Radici (in progress)



Datei Bearbeiten Ansicht Favoriten Extras ?
 T

Adresse http://www.pv.infn.it/~radici/FFdatabase/

**\*\*\* FRAGMENTATION FUNCTIONS \*\*\***

[sitemap](#)
[text on FF](#)
[references](#)
[parametrizations](#)
[links](#)
[ESOP](#)

---

**sitemap:**

**sitemap (this page)**

<p style="text-align: center;"> <u>text on fragmentation functions</u>            (../FFdatabase/text.html)         </p> <p><a href="#">parton distribution functions (PDFs)</a></p> <p><a href="#">unintegrated PDFs</a></p> <p><a href="#">fragmentation functions (FFs)</a></p> <p><a href="#">unintegrated FFs</a></p> <p><a href="#">multiple-hadron FFs</a></p> <p><a href="#">models calculations</a></p> <p><a href="#">target fragmentation and fracture functions</a></p>	<p style="text-align: center;"> <u>references related to fragmentation functions</u>            (../FFdatabase/references.html)         </p> <p><a href="#">operator definitions of PDFs and FFs</a></p> <p><a href="#">information on PDF's / parametrizations</a></p> <p><a href="#">information on FF's / parametrizations</a></p> <p><a href="#">models for FF's</a></p> <p><a href="#">evolution of FF's / scaling violations</a></p> <p><a href="#">target fragmentation and fracture functions</a></p> <p><a href="#">more references (still not properly sorted)</a></p>	<p style="text-align: center;"> <u>parametrizations</u>            (../FFdatabase/parametrizations.html)         </p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; padding: 5px;"> <p><a href="#">Stefan Kretzer</a></p> <p style="text-align: center;"></p> </td> <td style="width: 50%; padding: 5px;"> <p style="text-align: center;"></p> </td> </tr> <tr> <td style="padding: 5px;"> <p><a href="#">Kniehl, Kramer, Pötter</a></p> <p style="text-align: center;"></p> </td> <td style="padding: 5px;"> <p>all three combined in one <a href="#">FORTRAN library</a></p> </td> </tr> <tr> <td style="padding: 5px;"> <p><a href="#">Bourhis, Fontannaz, Guillet, Werlen</a> (soon to come)</p> </td> <td style="padding: 5px;"> <p>(courtesy of S.Kretzer)</p> </td> </tr> </table>	<p><a href="#">Stefan Kretzer</a></p> <p style="text-align: center;"></p>	<p style="text-align: center;"></p>	<p><a href="#">Kniehl, Kramer, Pötter</a></p> <p style="text-align: center;"></p>	<p>all three combined in one <a href="#">FORTRAN library</a></p>	<p><a href="#">Bourhis, Fontannaz, Guillet, Werlen</a> (soon to come)</p>	<p>(courtesy of S.Kretzer)</p>
<p><a href="#">Stefan Kretzer</a></p> <p style="text-align: center;"></p>	<p style="text-align: center;"></p>							
<p><a href="#">Kniehl, Kramer, Pötter</a></p> <p style="text-align: center;"></p>	<p>all three combined in one <a href="#">FORTRAN library</a></p>							
<p><a href="#">Bourhis, Fontannaz, Guillet, Werlen</a> (soon to come)</p>	<p>(courtesy of S.Kretzer)</p>							

[sitemap](#)  
[text on FF](#)  
[references](#)  
[parametrizations](#)  
  
[links](#)

this database is a project of the [ESOP](#) network

maintained by [Marco Radici](#) and [Rainer Jakob](#)

Last modified: Mon, Jan 28, 2002, 13:29



Datei Bearbeiten Ansicht Favoriten Extras ?

Adresse <http://www.pv.infn.it/~radici/FFdatabase/>

**\* FF \***

**\*\*\* FRAGMENTATION FUNCTIONS \*\*\***

[sitemap](#) [text on FF](#) [references](#) [parametrizations](#) [links](#) [ESOP](#)

[sitemap](#)

[text on FF](#)

[references](#)

[parametrizations](#)

[links](#)

[ESOP](#)

**sitemap:**

all this and much more ... Rainer Jakob & Marco Radici

**database on fragmentation functions**

systematics | references | **models** | parametrizations

unintegrated FFs

multiple-hadron FFs

models calculations

target fragmentation and fracture functions

models for FF's

evolution of FF's / scaling violations

target fragmentation and fracture functions

more references (still not properly sorted)

Bourhis, Fontannaz, Guillet, Werlen (soon to come)

library (courtesy of S.Kretzer)

maintained by [Marco Radici](#) and [Rainer Jakob](#)

Last modified: Mon, Jan 28, 2002, 13:29



<http://www.pv.infn.it/~radici/FFdatabase/>

all this and much more ...

Rainer Jakob & Marco Radici

## database on fragmentation functions

systematics | references | **models** | parametrizations

maintained by [Marco Radici](#) and [Rainer Jakob](#)

Last modified: Mon, Jan 28, 2002, 13:29

[unintegrated FFs](#)

[multiple-hadron FFs](#)

[models calculations](#)

[target fragmentation and fracture functions](#)

[models for FF's](#)

[evolution of FF's / scaling violations](#)

[target fragmentation and fracture functions](#)

[more references \(still not properly sorted\)](#)



[Bourhis, Fontannaz, Guillet, Werlen](#)  
(soon to come)

[library](#)

(courtesy of S.Kretzer)



# summary

## one hadron FF

	without $\mathbf{k}_T$	with $\mathbf{k}_T$
spin-0	$D_1$	$H_1^\perp$
spin-1/2	$D_1, G_1, H_1$	$D_{1T}^\perp, H_1^\perp, G_{1T}, H_{1L}^\perp, H_{1T}^\perp$
spin-1	$D_1, D_{1LL}, G_1, H_1, H_{1LT}$	$D_{1T}^\perp, H_1^\perp, G_{1T}, H_{1L}^\perp, H_{1T}^\perp,$ $D_{1T}^\perp, D_{1LT}, D_{1TT}, G_{1LT}, G_{1TT},$ $H_{1LL}^\perp, H'_{1LT}, H_{1LT}^\perp$

## two hadron FF

	without $\mathbf{k}_T$	with $\mathbf{k}_T$
spin-0	$D_1, H_1^\diamond$	$G_1^\perp, H_1^\perp$

higher twist



# conclusions

- the number of independent fragmentation functions is limited and there is a **simple systematics** behind
- **spin-dependent fragmentation functions** are not only a tool for the extraction of distribution functions, but provide the key information for understanding hadronic structure